

Abstract Submitted
for the DAMOP01 Meeting of
The American Physical Society

Sorting Category: 26 (Experimental)

2s Electron Loss to the Continuum from Metastable $C^{4+}(1s2s, ^3S)$ in C^{4+} Plus Helium Collisions¹ H. ALIABADI, , P. RICHARD, , C.P. BHALLA, J R Macdonald Laboratory, Kansas State University, Manhattan KS 66506, M. GEALY, Concordia College, Morehead, MN 56562 — We have investigated the broad peak at 1200eV in the electron production double differential cross section of 30MeV C^{4+} collisions with a helium target at various laboratory scattering angles. In the present experiment this peak is observed at angles between 15° and 160° . A simple model treating the 2s electron of the metastable projectile as elastically scattering from the helium atom predicts an electron loss peak in the observed cross sections. The model which includes the Compton profile of the 2s electron in $C^{4+}(1s2s, ^3S)$ predicts the observed cross sections when the metastable fraction is taken into account. The data and comparisons with the model calculations will be presented.

¹The Chemical Sciences, Geosciences and Biosciences Division, Office of Basic Energy Sciences, Office of Science, the U.S. Department of Energy supported this work

Prefer Oral Session

Prefer Poster Session

J R Macdonald Laboratory, Kansas State University, Manhattan KS 66506

Habibollah Aliabadi
habib@phys.ksu.edu

Date submitted: February 5, 2001

Electronic form version 1.4